

NECK JUNCTIONAL HEMORRHAGE CONTROL



01
EXPOSE the injury and assess the bleeding source.



02
APPLY DIRECT PRESSURE to bleeding source, use dressing from casualty's JFAK.



03
PACK wound tightly with hemostatic gauze until the wound cavity is filled.



04
ENSURE gauze extends 1-2 inches above the skin.



05
HOLD pressure for a minimum of 3 minutes.



06
REASSESS to ensure bleeding has been controlled while maintaining pressure.

IF BLEEDING HAS NOT BEEN CONTROLLED:

07a If packed with hemostatic dressing, remove and repack starting at **STEP 03**.

OR

07b If packed with gauze, apply additional gauze and pressure (for 3 min) until bleeding stops.



08
PLACE an elastic bandage over the dressing leaving enough tail to tie it into a knot.



09
While maintaining pressure, **WRAP** bandage (no less than 1½ times) over the packing material covering it completely.



WRAP diagonally across the chest under the opposite arm (armpit) around the back to the neck and back over the wound.

Continued on the next page...

NECK JUNCTIONAL HEMORRHAGE CONTROL

Continued...




10 **WRAP** around neck and under the arm (on alternating sides of the tail, while maintaining tension/pressure), pulling elastic bandage tightly for pressure, covering the packing material.

NOTE: Do *not* use pressure bar on the neck.



11 **SECURE** the dressing by tying a non-slip knot with end of elastic bandage and its tail.




12 **SECURE** elastic bandage tails with tape, wrapping a minimum of 1½ times around the knot.



13 **SWATH** the upper arm (of the injured side) to the chest using a bandage.



14 **CONTINUE TO ASSESS** wound for further bleeding.



15 **DOCUMENT** all findings and treatments on the DD1380 TCCC, Casualty Card.